



# B.I.O.Tox® Activ8

**Identification:**

Mineral feed

**Product description:**

Mineral feed containing substances for reduction of contamination of feed by mycotoxins

**Animal category:**

Ruminants, poultry and pigs

**Ingredients:**

Calcium:	0.9 %	Lysine:	0.0 %
Phosphorus:	0.0 %	Methionine:	0.0 %
Sodium:	1.3 %	Ash insoluble in HCl:	62.7 %
Magnesium:	2.9 %	Crude ash:	77.7 %
Moisture:	12.6 %		

**Additives/kg:***Binders:*

Sepiolite (1g562) 162 000 mg

*Mycotoxin reducers:*

Bentonite (1m558) 720 000 mg

*Flavouring compounds:*

Dry grape extract of Vitis vinifera spp. vinifera (2b485) 1 000 mg

**Composition:**

5.34.1 Thistle seed (ground), 7.13.1 Vegetal carbon (charcoal)

**Recommended dosage:**

Low content of mycotoxins / preventive: 0.5 – 1.0 kg/t

Increased content of mycotoxins: 1.0 – 2.0 kg/t

Required dosage depends on several parameters including status of feed contamination, feed quality, animal species, animal age, health status, etc. In order to obtain maximum efficacy, mix B.I.O.Tox® Activ8 thoroughly while loading raw materials and/or feeds.

**Characteristics:**

Grey powder

Light variations in color do not affect product quality.

**Feeding advice:**

The simultaneous oral use with macrolides shall be avoided. For poultry the simultaneous use of robenidine shall be avoided as well. Due to the increased content of binders this mineral feed should only be applied up to 1.0% of the daily ration.

**Shelf life:**

24 months after manufacturing date when stored under dry conditions in original packaging. Store in a well-ventilated area.

**Package size:**

25 kg bag

Version: 09

Issued: 05/2025